



Jazz Pharmaceuticals Announces Ziihera® (zanidatamab-hrii) Investor Webcast on December 11, 2024

December 04, 2024

DUBLIN, Dec. 4, 2024 /PRNewswire/ -- Jazz Pharmaceuticals plc (Nasdaq: JAZZ) today announced that it will host a webcast on Wednesday, December 11, 2024, at 4:30 p.m. ET / 9:30 p.m. GMT to provide an overview of clinical data, patient need and commercialization strategy for Ziihera® (zanidatamab-hrii), the first chemotherapy-free dual HER2-targeted bispecific antibody indicated for biliary tract cancer (BTC). Ziihera was approved under accelerated approval by the U.S. Food and Drug Administration (FDA) on November 20, 2024, for the treatment of adults with previously treated, unresectable or metastatic HER2-positive (IHC 3+) BTC, as detected by an FDA-approved test.¹

Jazz senior management will provide an overview of Ziihera and commercial launch plans, and Dr. Shubham Pant will discuss previously disclosed BTC data from the HERIZON-BTC-01 trial. Shubham Pant, M.D., MBBS, is a professor in the Department of Gastrointestinal Medical Oncology with a joint appointment in the Department of Investigational Cancer Therapeutics at The University of Texas MD Anderson Cancer Center.

Audio webcast/conference call:

U.S. Dial-In Number: +1 800 715 9871

Ireland Dial-In Number: +353 1800 943 926

Additional global dial-in numbers are available [here](#).

Passcode: 4898380

Interested parties may access the live audio webcast via the Investors section of the Jazz Pharmaceuticals website at <https://investor.jazzpharma.com/investors/events-presentations>. To ensure a timely connection, it is recommended that participants register at least 15 minutes prior to the scheduled webcast.

A replay of the webcast will be available via the Investors section of the Jazz Pharmaceuticals website at <https://investor.jazzpharma.com/investors/events-presentations>.

About Ziihera® (zanidatamab-hrii)

Ziihera (zanidatamab-hrii) is a bispecific HER2-directed antibody that binds to two extracellular sites on HER2. Binding of zanidatamab-hrii with HER2 results in internalization leading to a reduction of the receptor on the tumor cell surface. Zanidatamab-hrii induces complement-dependent cytotoxicity (CDC), antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP). These mechanisms result in tumor growth inhibition and cell death in vitro and in vivo.¹ In the United States, Ziihera is indicated for the treatment of adults with previously treated, unresectable or metastatic HER2-positive (IHC 3+) biliary tract cancer (BTC), as detected by an FDA-approved test.¹ The U.S. Food and Drug Administration (FDA) granted accelerated approval for this indication based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).¹

Zanidatamab is not approved anywhere else in the world.

Zanidatamab is being developed in multiple clinical trials as a targeted treatment option for patients with solid tumors that express HER2. Zanidatamab is being developed by Jazz and BeiGene, Ltd. (BeiGene) under license agreements from Zymeworks, which first developed the molecule.

The FDA granted Breakthrough Therapy designation for zanidatamab development in patients with previously treated HER2 gene-amplified BTC, and two Fast Track designations for zanidatamab: one as a single agent for refractory BTC and one in combination with standard-of-care chemotherapy for 1L gastroesophageal adenocarcinoma (GEA). Additionally, zanidatamab has received Orphan Drug designations from FDA for the treatment of BTC and GEA, as well as Orphan Drug designation from the European Medicines Agency for the treatment of BTC and gastric cancer.

Important Safety Information

WARNING: EMBRYO-FETAL TOXICITY

Exposure to ZIIHERA during pregnancy can cause embryo-fetal harm. Advise patients of the risk and need for effective contraception.

WARNINGS AND PRECAUTIONS

Embryo-Fetal Toxicity

ZIIHERA can cause fetal harm when administered to a pregnant woman. In literature reports, use of a HER2-directed antibody during pregnancy resulted in cases of oligohydramnios and oligohydramnios sequence manifesting as pulmonary hypoplasia, skeletal abnormalities, and neonatal death.

Verify the pregnancy status of females of reproductive potential prior to the initiation of ZIIHERA. Advise pregnant women and females of reproductive potential that exposure to ZIIHERA during pregnancy or within 4 months prior to conception can result in fetal harm. Advise females of reproductive potential to use effective contraception during treatment with ZIIHERA and for 4 months following the last dose of ZIIHERA.

Left Ventricular Dysfunction

ZIIHERA can cause decreases in left ventricular ejection fraction (LVEF). LVEF declined by >10% and decreased to <50% in 4.3% of 233 patients. Left ventricular dysfunction (LVD) leading to permanent discontinuation of ZIIHERA was reported in 0.9% of patients. The median time to first occurrence

of LVD was 5.6 months (range: 1.6 to 18.7). LVD resolved in 70% of patients.

Assess LVEF prior to initiation of ZIIHERA and at regular intervals during treatment. Withhold dose or permanently discontinue ZIIHERA based on severity of adverse reactions.

The safety of ZIIHERA has not been established in patients with a baseline ejection fraction that is below 50%.

Infusion-Related Reactions

ZIIHERA can cause infusion-related reactions (IRRs). An IRR was reported in 31% of 233 patients treated with ZIIHERA as a single agent in clinical studies, including Grade 3 (0.4%), and Grade 2 (25%). IRRs leading to permanent discontinuation of ZIIHERA were reported in 0.4% of patients. IRRs occurred on the first day of dosing in 28% of patients; 97% of IRRs resolved within one day.

Prior to each dose of ZIIHERA, administer premedications to prevent potential IRRs. Monitor patients for signs and symptoms of IRR during ZIIHERA administration and as clinically indicated after completion of infusion. Have medications and emergency equipment to treat IRRs available for immediate use.

If an IRR occurs, slow, or stop the infusion, and administer appropriate medical management. Monitor patients until complete resolution of signs and symptoms before resuming. Permanently discontinue ZIIHERA in patients with recurrent severe or life-threatening IRRs.

Diarrhea

ZIIHERA can cause severe diarrhea.

Diarrhea was reported in 48% of 233 patients treated in clinical studies, including Grade 3 (6%) and Grade 2 (17%). If diarrhea occurs, administer antidiarrheal treatment as clinically indicated. Perform diagnostic tests as clinically indicated to exclude other causes of diarrhea. Withhold or permanently discontinue ZIIHERA based on severity.

ADVERSE REACTIONS

Serious adverse reactions occurred in 53% of 80 patients with unresectable or metastatic HER2-positive BTC who received ZIIHERA. Serious adverse reactions in >2% of patients included biliary obstruction (15%), biliary tract infection (8%), sepsis (8%), pneumonia (5%), diarrhea (3.8%), gastric obstruction (3.8%), and fatigue (2.5%). A fatal adverse reaction of hepatic failure occurred in one patient who received ZIIHERA.

The most common adverse reactions in 80 patients with unresectable or metastatic HER2-positive BTC who received ZIIHERA (≥20%) were diarrhea (50%), infusion-related reaction (35%), abdominal pain (29%), and fatigue (24%).

USE IN SPECIFIC POPULATIONS

Pediatric Use

Safety and efficacy of ZIIHERA have not been established in pediatric patients.

Geriatric Use

Of the 80 patients who received ZIIHERA for unresectable or metastatic HER2-positive BTC, there were 39 (49%) patients 65 years of age and older. Thirty-seven (46%) were aged 65-74 years old and 2 (3%) were aged 75 years or older.

No overall differences in safety or efficacy were observed between these patients and younger adult patients.

About Jazz Pharmaceuticals

Jazz Pharmaceuticals plc (Nasdaq: JAZZ) is a global biopharma company whose purpose is to innovate to transform the lives of patients and their families. We are dedicated to developing life-changing medicines for people with serious diseases — often with limited or no therapeutic options. We have a diverse portfolio of marketed medicines, including leading therapies for sleep disorders and epilepsy, and a growing portfolio of cancer treatments. Our patient-focused and science-driven approach powers pioneering research and development advancements across our robust pipeline of innovative therapeutics in oncology and neuroscience. Jazz is headquartered in Dublin, Ireland with research and development laboratories, manufacturing facilities and employees in multiple countries committed to serving patients worldwide. Please visit www.jazzpharmaceuticals.com for more information.

Contacts:

Investors:

Jeff Macdonald
Executive Director, Investor Relations
Jazz Pharmaceuticals plc
InvestorInfo@jazzpharma.com
Ireland +353 1 634 3211
U.S. +1 650 496 2717

Media:

Kristin Bhavnani
Head of Global Corporate Communications
Jazz Pharmaceuticals plc
CorporateAffairsMediaInfo@jazzpharma.com
Ireland +353 1 637 2141
U.S. +1 215 867 4948

References:

¹ZIIHERA (zanidatamab-hrii) Prescribing Information. Palo Alto, CA: Jazz Pharmaceuticals, Inc.



View original content to download multimedia: <https://www.prnewswire.com/news-releases/jazz-pharmaceuticals-announces-ziihera-zanidatamab-hrii-investor-webcast-on-december-11-2024-302323036.html>

SOURCE Jazz Pharmaceuticals plc