



## Journal of Clinical Oncology publishes pivotal Phase 3 data for Jazz Pharmaceuticals' Vyxeos® (daunorubicin and cytarabine) Liposome for Injection

July 19, 2018

DUBLIN, July 19, 2018 /PRNewswire/ -- Jazz Pharmaceuticals plc (Nasdaq: JAZZ) announced today that data from the pivotal Phase 3 study of Vyxeos® (daunorubicin and cytarabine) liposome for injection compared to standard of care cytarabine and daunorubicin (7+3) were published online in the [Journal of Clinical Oncology](#). The study evaluated the efficacy and safety of Vyxeos compared to 7+3 in 309 patients who were between 60 to 75 years of age with newly diagnosed therapy-related acute myeloid leukemia (t-AML) or AML with myelodysplasia-related changes (AML-MRC), a rapidly progressing and life-threatening blood cancer.

The study met its primary endpoint as Vyxeos demonstrated a superior improvement in overall survival compared to the 7+3 treatment regimen. The median overall survival for the Vyxeos treatment group was 9.6 months compared with 5.9 months for the 7+3 treatment group (2-sided p value = 0.005; HR [95% confidence interval] = 0.69 [0.52, 0.90]). Vyxeos was also associated with a significantly higher remission rate than 7+3 with a complete response rate of 38% versus 26%; p=0.036. In addition, the overall rate of hematopoietic stem cell transplant (HSCT) was 34% in the Vyxeos arm and 25% in the 7+3 arm. The reported adverse reactions with Vyxeos were generally consistent with the known safety profile of cytarabine and daunorubicin therapy.

Vyxeos was approved by the U.S. Food and Drug Administration (FDA) in August 2017, and is the first FDA-approved treatment specifically indicated for adults with newly-diagnosed t-AML or AML-MRC. Data from the study formed the basis of the FDA application and the Marketing Authorization Application (MAA) to the European Medicines Agency's Committee for Medicinal Products for Human Use (CHMP).

"We are encouraged by the positive response to Vyxeos from U.S. health care professionals who had been waiting for an advancement in the treatment of these two types of patients with AML," said Allen Yang, M.D., Ph.D., vice president and acting chief medical officer of Jazz Pharmaceuticals. "At Jazz we are keenly aware that every clinical trial result advances the science to help patients and we are committed to helping as many people as possible with Vyxeos."

Designed with Jazz's CombiPlex® proprietary technology, Vyxeos is a unique liposomal formulation that delivers a fixed-ratio of daunorubicin and cytarabine to the bone marrow that has been shown to have synergistic effects at killing leukemia cells in vitro and in animal models.

"Vyxeos is the first agent to significantly improve survival in older, fit AML patients with secondary AML," said Jeffrey E. Lancet, MD, Chair of the Department of Malignant Hematology at Moffitt Cancer Center and lead author of the publication. "Collectively, the Phase 3 clinical data support the adoption of Vyxeos for the treatment of adults with newly-diagnosed t-AML or AML-MRC."

**Vyxeos has different dosage recommendations from other medications that contain daunorubicin and/or cytarabine. Do not substitute Vyxeos for other daunorubicin- and/or cytarabine- containing products.**

In the Phase 3 study, patients in the Vyxeos arm received 44mg/100mg per m<sup>2</sup> (daunorubicin and cytarabine) liposome intravenously via a 90 minute infusion on days 1, 3 and 5 of induction (days 1 and 3 if a second induction was needed) and 29mg/65mg per m<sup>2</sup> (daunorubicin and cytarabine) liposome on days 1 and 3 for consolidation. Patients in the 7+3 arm received induction with cytarabine 100mg/m<sup>2</sup>/day on days 1-7 by continuous infusion and daunorubicin 60mg/m<sup>2</sup>/day on days 1-3. For consolidation, cytarabine was dosed on days 1-5 and daunorubicin on days 1-2. Patients could receive up to 2 cycles of induction and 2 cycles of consolidation in each arm. Subsequent induction was recommended for patients who did not achieve a response and was mandatory for patients achieving >50% reduction in percent blasts.

For the primary endpoint of overall survival, Vyxeos demonstrated an improvement that was superior to the 7+3 treatment regimen. The median overall survival for the Vyxeos treatment group was 9.6 months compared with 5.9 months for the 7+3 treatment group (2-sided p value = 0.005; HR [95% confidence interval] = 0.69 [0.52, 0.90]). Vyxeos also was associated with a significantly higher remission rate than 7+3 with a complete response rate of 38% versus 26%; p=0.036. In addition, the overall rate of hematopoietic stem cell transplant (HSCT) was 34% in the Vyxeos arm and 25% in the 7+3 arm. The overall, all-cause 30-day mortality was 6% in the Vyxeos arm and 11% in the 7+3 arm. During the first 60 days of the study, 14% (21/153) of patients died in the Vyxeos arm vs. 21% (32/151) of patients in the 7+3 treatment group.

The incidences of nonhematologic adverse events were comparable between arms, despite a longer treatment phase and prolonged time to neutrophil and platelet count recovery with Vyxeos. Fatal treatment-emergent CNS hemorrhage not in the setting of progressive disease occurred in 2% of patients in the Vyxeos arm and 0.7% of patients in the control arm. Six percent of patients in both the Vyxeos and control arm had a fatal adverse reaction on treatment or within 30 days of therapy that was not in the setting of progressive disease. The most common adverse reactions (incidence ≥ 25%) were bleeding events, fever, rash, swelling, nausea, sores in the mouth or throat, diarrhea, constipation, muscle pain, tiredness, stomach pain, difficulty breathing, headache, cough, decreased appetite, irregular heartbeat, pneumonia, blood infection, chills, sleep disorders and vomiting.

### About Vyxeos®

Vyxeos® (daunorubicin and cytarabine) liposome for injection 44mg/100mg is a liposome formulation of a fixed combination of daunorubicin and cytarabine for intravenous infusion.<sup>1</sup> Vyxeos is indicated for the treatment of adults with newly-diagnosed t-AML or AML-MRC. For more information about Vyxeos in the United States, please visit <https://vyxeos.com>.

### Important Safety Information

**Vyxeos has different dosage recommendations from other medications that contain daunorubicin and/or cytarabine. Do not substitute Vyxeos for other daunorubicin- and/or cytarabine- containing products.**

Vyxeos should not be given to patients who have a history of serious allergic reaction to daunorubicin, cytarabine or any of its ingredients.

Vyxeos can cause a severe decrease in blood cells (red and white blood cells and cells that prevent bleeding, called platelets) which can result in serious infection or bleeding and possibly lead to death. Your doctor will monitor your blood counts during treatment with Vyxeos. Patients should tell the doctor about new onset fever or symptoms of infection or if they notice signs of bruising or bleeding.

Vyxeos can cause heart-related side effects. Tell your doctor about any history of heart disease, radiation to the chest, or previous chemotherapy. Inform your doctor if you develop symptoms of heart failure such as:

- shortness of breath or trouble breathing
- swelling or fluid retention, especially in the feet, ankles or legs
- unusual tiredness

Vyxeos may cause allergic reactions including anaphylaxis. Seek immediate medical attention if you develop signs and symptoms of anaphylaxis such as:

- trouble breathing
- severe itching
- skin rash or hives
- swelling of the face, lips, mouth, or tongue

Vyxeos contains copper and may cause copper overload in patients with Wilson's disease or other copper-processing disorders.

Vyxeos can damage the skin if it leaks out of the vein. Tell your doctor right away if you experience symptoms of burning, stinging, or blisters and skin sores at the injection site.

Vyxeos can harm your unborn baby. Inform your doctor if you are pregnant, planning to become pregnant, or nursing. Do not breastfeed while receiving Vyxeos. Females and males of reproductive potential should use effective contraception during treatment and for 6 months following the last dose of Vyxeos.

The most common side effects were bleeding events, fever, rash, swelling, nausea, sores in the mouth or throat, diarrhea, constipation, muscle pain, tiredness, stomach pain, difficulty breathing, headache, cough, decreased appetite, irregular heartbeat, pneumonia, blood infection, chills, sleep disorders, and vomiting.

Please see full Prescribing Information for Vyxeos including BOXED Warning at: <http://pp.jazzpharma.com/pi/vyxeos.en.USPI.pdf>

#### **About AML**

Acute myeloid leukemia (AML) is a blood cancer that begins in the bone marrow, which produces most of the body's new blood cells.<sup>2</sup> AML cells crowd out healthy cells and move aggressively into the bloodstream to spread cancer to other parts of the body.<sup>3</sup> AML is a relatively rare disease representing 1.3 percent of all new cancer cases.<sup>4</sup> It is estimated that more than 19,500 people will be diagnosed with AML in the United States this year with the potential for nearly 11,000 people to die from the disease.<sup>5</sup> The median age at diagnosis is 68 years old,<sup>4</sup> with rising age associated with a progressively worsening prognosis.<sup>6</sup> There is also a reduced tolerance for intensive chemotherapy as patients age.<sup>7</sup> AML has the lowest survival rate of any other form of leukemia.<sup>4</sup> Patients with newly diagnosed t-AML or AML-MRC may have a particularly poor prognosis.<sup>8-10</sup> A hematopoietic stem cell transplant (HSCT) may be a curative treatment option for patients.<sup>11</sup>

#### **About Jazz Pharmaceuticals plc**

Jazz Pharmaceuticals plc (Nasdaq: JAZZ) is an international biopharmaceutical company focused on improving patients' lives by identifying, developing and commercializing meaningful products that address unmet medical needs. The company has a diverse portfolio of products and product candidates with a focus in the areas of sleep and hematology/oncology. In these areas, Jazz Pharmaceuticals markets Xyrem® (sodium oxybate) oral solution, Erwinaze® (asparaginase *Erwinia chrysanthemi*), Defitelio® (defibrotide sodium) and Vyxeos® (daunorubicin and cytarabine) liposome for injection in the U.S. and markets Erwinaze® and Defitelio® (defibrotide) in countries outside the U.S. For country-specific product information, please visit [www.jazzpharmaceuticals.com/products](http://www.jazzpharmaceuticals.com/products). For more information, please visit [www.jazzpharmaceuticals.com](http://www.jazzpharmaceuticals.com) and follow us on Twitter at [@JazzPharma](https://twitter.com/JazzPharma).

#### **References:**

1. Vyxeos [package insert]. Palo Alto, CA: Jazz Pharmaceuticals; 2017.
2. National Cancer Institute. General Information About Adult Acute Myeloid Leukemia <https://www.cancer.gov/types/leukemia/patient/adult-aml-treatment-pdq> Accessed June 7, 2017.
3. American Cancer Society. What is Acute Myeloid Leukemia? <https://www.cancer.org/cancer/acute-myeloid-leukemia/about/what-is-aml.html> Accessed March 20, 2017.
4. SEER Stat Facts: AML. 2017.
5. American Cancer Society. Key Statistics About Acute Myeloid Leukemia. <https://www.cancer.org/cancer/acute-myeloid-leukemia/about/key-statistics.html> Accessed May 24, 2018.
6. Baer MR, et al., Leukemia, 2011 May; 25(5):10.1038/eu.2011.9.
7. Klepin HD. Hematology Am Soc Hematol Educ Program. 2014;2014(1):8-13.
8. Goldstone AH, Burnett AK, Avivi I et al. Secondary acute myeloid leukemia has a worse outcome than de novo AML, even taking into account cytogenetics and age. AML 10, 11, 12 MRC Trials. Blood 2002; 100 (88a): (Abstr 322).
9. Schiller GJ, Hematol Educ Program, 2013:201-208.

10. Kern W, Haferlach T, Schnittger S, Hiddemann W, Schoch C. Prognosis in therapy-related acute myeloid leukemia and impact of karyotype. [J Clin Oncol](#). 2004 Jun 15;22(12):2510-1.
11. Peccatori, J and Ciceri, F. *Haematologica*. 2010 Jun; 95(6): 857–859. doi: 10.3324/haematol.2010.023184.



View original content with multimedia: <http://www.prnewswire.com/news-releases/journal-of-clinical-oncology-publishes-pivotal-phase-3-data-for-jazz-pharmaceuticals-vyxeos-daunorubicin-and-cytarabine-liposome-for-injection-300684026.html>

SOURCE Jazz Pharmaceuticals plc

Jacqueline Kirby, Vice President, Corporate Affairs & Government Relations, Ireland +353 1 697 2141, U.S. +1 215 867 4910, Investor Contact:  
Kathee Littrell, Vice President, Investor Relations, Ireland +353 1 634 7887, U.S. +1 650 496 2717